



TRANSFORMING
AND ANALYZING
RATIONAL
FUNCTIONS

Rational Functions

- A rational function is a function composed of a ratio of two polynomial functions, $p(x)$ and $q(x)$.

$$\circ r(x) = \frac{p(x)}{q(x)}$$

- For a rational function, the general form is $f(x) = \frac{a}{bx-c} + d$, where a , b , c , and d are real numbers.
- The rational parent function is $f(x) = \frac{1}{x}$. This is also called an inverse variation function.
- The full family of rational functions is generated by applying transformations to the Rational parent function
- Transformations are applied using parameters that are multiplied or added to the independent variable in the functional relationship

Changes in a

- The parameter a influences the vertical stretch or compression of the graph of the parabola.
- If $|a| > 1$, then the y -values are multiplied by a factor of a to vertically stretch the graph
- If $0 < |a| < 1$, then the y -values are multiplied by a factor of a to vertically compress the graph
- If $a < 0$, then the graph will be reflected across the x -axis

Changes in b

- The parameter b influences the horizontal stretch or compression of the graph of the parabola.
- If $|b| > 1$, then the x-values are multiplied by a factor of $\frac{1}{|b|}$ to horizontally compress the graph
- If $0 < |b| < 1$, then the x-values are multiplied by a factor of $\frac{1}{|b|}$ to horizontally stretch the graph
- If $b < 0$, then the graph will be reflected across the y-axis

Changes in c

- The parameter c , like b , influences the horizontal translation of the graph of the parabola.
- Note that in the general form, the sign in front of the c is negative. This means that when reading the value of c from the equation, you should read the opposite sign from what is given in the equation.
- If $c > 0$, then the graph will translate $\left| \frac{c}{b} \right|$ to the right.
- If $c < 0$, then the graph will translate $\left| \frac{c}{b} \right|$ to the left.

Changes in d

- The parameter d influences the vertical translation of the graph of the parabola.
- If $d > 0$, then the graph of the parabola will translate $|d|$ units up.
- If $d < 0$, then the graph of the parabola will translate $|d|$ units down.

Asymptotes

- Each rational function has two asymptotes: one horizontal and one vertical
- The horizontal asymptote is governed by vertical parameters changes. A vertical translation moves the asymptote d units and a vertical dilation does not move the asymptote.
- The vertical asymptote is governed by horizontal parameters changes. A horizontal translation moves the asymptote c units and a horizontal dilation moves the asymptote closer or away from the x -axis by a factor of $\frac{1}{b}$.
 - horizontal asymptote: $y = d$
 - vertical asymptote: $x = \frac{c}{b}$

Domain and Range

- A rational function involves the ratio of two polynomial functions. Since the function in the denominator can never equal 0, any values of x that cause the denominator to equal 0 are excluded from the domain. Therefore, the domain will always be *all real numbers minus $\frac{c}{b}$* , or $\{x \mid x \in \mathbb{R}, x \neq \frac{c}{b}\}$
- The range is restricted by the horizontal asymptote, $y = d$. Therefore, $y = d$ must be excluded from the range of a rational function. $\{y \mid y \in \mathbb{R}, y \neq d\}$

X- and Y-intercepts

- A rational function has at most one x-intercepts. If it exists, the x-intercept is located at:

- $(\frac{cd - a}{bd}, 0)$

- The values of b and d may not equal 0. When $b = 0$, the function does not exist since there is no x-term in the function. When $d = 0$, the x-axis is an asymptote and the x-intercept does not exist.

- A rational function has at most one y-intercepts. If it exists, the y-intercept is located at:

- $(0, \frac{a}{-c} + d)$

- When $c = 0$, the y-axis is an asymptote and they-intercept does not exist.

Examples

- What transformations of the rational parent function, $f(x) = \frac{1}{x}$, will result in the graph of the rational function $g(x) = -\frac{3}{x-2} + 1.5$?

Examples

- Step 1: Rewrite the equation of $g(x)$ in general form $y = \frac{a}{bx-c} + d$ to determine the values of the parameters a , b , c , and d .

- $g(x) = \frac{a}{bx-c} + d$

- $g(x) = -\frac{3}{x-2} + 1.5$

- $g(x) = \frac{-3}{1x-2} + 1.5$

- $a = -3, b = 1, c = 2, d = 1.5$

Examples

- Step 2: Use the parameters to describe the transformations of the rational parent function $f(x)$ that are necessary to produce $g(x)$.
- $a = -3$, so there is vertical stretch by a factor of 3; a is negative, so it is reflected over the x -axis
- $b = 1$, there is no change to the graph
- $c = 2$, so there is a horizontal shift $\frac{2}{1} = 2$ units to the right
- $d = 1.5$, so there is a vertical shift 1.5 units up

Examples

- Identify the key attributes of $y = -\frac{12}{3x} + 2$, including domain and range (including asymptotes), x- and y-intercepts. Write the domain and range in set builder notation.

Examples

- Step 1: Determine the domain and range of $y = -\frac{12}{3x} + 2$
 - The domain excludes $\frac{c}{b}$
 - $\frac{c}{b} = \frac{0}{3} = 0$
 - $\{x \mid x \in \mathbb{R}, x \neq 0\}$
 - The range excludes d
 - $d = 2$
 - $\{y \mid y \in \mathbb{R}, y \neq 2\}$

Examples

- Step 2: Determine if the function has an x-intercept.
- Use the calculator to determine the x-intercept.
 - x-int: $(-2, 0)$
- Step 3: Determine if the function has a y-intercept.
- Use the calculator to determine the y-intercept.
 - y-int: none