


# Tangents


# Tangent

- A tangent is a line in the same plane as a circle that intersects the circle in exactly one point, called the point of tangency.
- A common tangent is a line, ray, or segment that is tangent to two circles in the same plane.




# Tangent Theorem

- In a plane, a line is tangent to a circle if and only if it is perpendicular to a radius drawn to the point to tangency.



# Examples

- Determine whether  $\overline{GH}$  is tangent to  $\odot F$ . Justify your answer.



$$A^2 + B^2 = C^2$$
$$6^2 + 8^2 \neq 12^2$$


$$36 + 64 = 324$$

$$100 \neq 324$$

# Examples


- Determine whether GH is tangent to  $\odot F$ . Justify your answer.

- $6^2 + 8^2 = (6 + 12)^2$
- $36 + 64 = 18^2$
- $100 = 324$
- Not tangent



# Examples

- JH is tangent to  $\odot G$  at J. Find x



$$x^2 + 12^2 = (x+8)^2$$

$$\cancel{x^2} + 144 = \cancel{x^2} + \underline{16x} + 64$$

$$144 = 16x + 64$$

$$-64 \quad \quad -64$$

$$\frac{80}{16} = \frac{16x}{16}$$


$$(x+8)(x+8)$$

$$\underline{x^2 + 8x + 8x + 64}$$

# Examples

- JH is tangent to  $\odot G$  at J. Find x.


- $x^2 + 12^2 = (x + 8)^2$
- $x^2 + 144 = x^2 + 16x + 64$
- $80 = 16x$
- $5 = x$
- 5, 12, 13 is a Pythagorean Triple
- Tangent



)

# Examples

- QS is tangent to  $\odot R$  at Q. Find x.



$$a^2 + b^2 = c^2$$

$$4^2 + x^2 = (x+2)^2$$

$$16x^2 = x^2 + 4x + 4$$

$$16 = 4x + 4$$


$$12 = 4x$$

$$x = 3$$

# Examples


- QS is tangent to  $\odot R$  at Q. Find x.

- $x^2 + 4^2 = (x + 2)^2$
- $x^2 + 16 = x^2 + 4x + 4$
- $16 = 4x + 4$
- $12 = 4x$
- $3 = x$




# Congruent Tangents

- If two segments from the same exterior point are tangent to a circle, then they are congruent.



# Examples

- $AB$  and  $CB$  are tangent to  $\odot D$ . Find the value of  $x$ .



$$x + 15 = 2x - 5$$

# Examples

- $AB$  and  $CB$  are tangent to  $\odot D$ . Find the value of  $x$ .


- $x + 15 = 2x - 5$

- $15 = x - 5$

- $20 = x$

Subtract  $x$  from both sides


Add 5 to both sides



# Examples


- Find the value of  $x$ . Assume that segments that appear to be tangent are tangent.

$$3x + 8 = 26$$



# Examples

- Find the value of  $x$ . Assume that segments that appear to be tangent are tangent.



- $3x + 8 = 26$

- $3x = 18$


Subtract 8 from both sides

- $x = 6$


Divide by 3 on both sides

# Circumscribed Polygons

- When a polygon is circumscribed about a circle, all of the sides of the polygon are tangent to the circle.



Polygons are circumscribed.




Polygons are *not* circumscribed.

# Examples

- Find the perimeter of each polygon for the given information. Assume that segments that appear to be tangent are tangent.

- $QT = 4$ ,  $PT = 9$ ,  $SR = 13$


$$4 + 4 + 9 + 9 + 13 + 13$$



# Examples

- Find the perimeter of each polygon for the given information. Assume that segments that appear to be tangent are tangent.

- $QT = 4$ ,  $PT = 9$ ,  $SR = 13$




- $QT = 4$ ,  $QU = 4$ ;  $PT = 9$ ,  $PS = 9$ ;  $RS = 13$ ,  $RU = 13$

- $\text{Perimeter} = 4 + 4 + 9 + 9 + 13 + 13 = 52$


# Examples

- Find the perimeter of each polygon for the given information. Assume that segments that appear to be tangent are tangent.
- $H I J K$  is a rhombus,  $S I = 5$ ,  $H R = 13$



# Examples

- Find the perimeter of each polygon for the given information. Assume that segments that appear to be tangent are tangent.



- $HIJK$  is a rhombus,  $SI = 5$ ,  $HR = 13$
- $SI = 5$ ,  $TI = 5$ ,  $RK = 5$ ,  $UK = 5$ ;  $HR = 13$ ,  $HS = 13$ ,  $JU = 13$ ,  $JT = 13$
- $Perimeter = 5 + 5 + 5 + 5 + 13 + 13 + 13 + 13 = 72$